

IE 400 Principles of Engineering Management

The Simplex Algorithm-II: Set 4

The Simplex Algorithm:

- To begin the simplex algorithm, convert the LP into a standard form,
- Convert the objective function $z = c_1x_1 + c_2x_2 + \dots + c_nx_n$ to the row-0 format:

$$z - c_1x_1 - c_2x_2 - \dots - c_nx_n = 0$$

The Simplex Algorithm:

$$\max \quad z = x_1 + 3x_2$$

s.t.

$$x_1 + x_2 \leq 6 \quad (1)$$

$$-x_1 + 2x_2 \leq 8 \quad (2)$$

$$x_1, x_2 \geq 0$$



$$\max \quad z$$

s.t.

$$z - x_1 - 3x_2 = 0 \quad (0)$$

$$x_1 + x_2 + s_1 = 6 \quad (1)$$

$$-x_1 + 2x_2 + s_2 = 8 \quad (2)$$

$$x_1, x_2, s_1, s_2 \geq 0$$

The Simplex Algorithm

Canonical Form 0:

Row	Basic Variable									RHS		
0		z	-	x_1	-	$3x_2$				=	0	
1				x_1	+	x_2	+	s_1		=	6	
2			-	x_1	+	$2x_2$			+	s_2	=	8

A system of linear equations in which each equation has a variable with a coefficient 1 in that equation (and a zero coefficient in all other equations) is said to be in *canonical form*.

If the RHS of each constraint in a canonical form is nonnegative, a basic feasible solution can be obtained by inspection.

The Simplex Algorithm

Recall that the Simplex Algorithm begins with an initial bfs and attempts to find better ones. After obtaining a canonical form, we search for the **initial bfs**.

By inspection, if we set $x_1=x_2=0$, we can solve for the values of s_1 and s_2 by setting s_i equal to the RHS of row i .

$$BV=\{s_1,s_2\} \text{ and } NBV=\{x_1,x_2\}$$

The Simplex Algorithm

You may also verify the calculations for the initial basic feasible solution by:

$$B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad x_N = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \quad x_B = \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}$$

The columns are
linearly independent

Setting $x_N = 0$ and solving for x_B : $Bx_B = b \Rightarrow x_B = B^{-1}b$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} = \begin{pmatrix} 6 \\ 8 \end{pmatrix} \Rightarrow \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}^{-1} \begin{pmatrix} 6 \\ 8 \end{pmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{pmatrix} 6 \\ 8 \end{pmatrix} = \begin{pmatrix} 6 \\ 8 \end{pmatrix}$$

The Simplex Algorithm

Notice that each basic variable is associated with the row of the canonical form in which the basic variable has a coefficient of 1.

To perform the simplex algorithm, we also need a basic variable **(not necessarily nonnegative!)** for row 0.

Observe that variable z appears in row 0 with a coefficient of 1, and z does not appear in any other row. Therefore, **we use z as basic variable for row 0.**

The Simplex Algorithm

Let us denote the initial canonical form as **canonical form 0**.

With this convention, the basic and nonbasic variables for the canonical form 0 are $BV=\{z,s_1,s_2\}$ and $NBV=\{x_1,x_2\}$.

For this basic feasible solution, $z=0$, $s_1=6$, $s_2=8$, $x_1=0$, $x_2=0$.

The Simplex Algorithm

In summary, the **canonical form**:

- LP has equality constraints and nonnegativity constraints ,
- There is one **basic** variable for each equality constraint,
- The column for the basic variable for constraint i has a 1 in constraint i and 0's elsewhere,
- The remaining variables are called **nonbasic**.

The Simplex Algorithm

Canonical Form 0:

Row	Basic Variable									RHS
0		z	-	x_1	-	$3x_2$				= 0
1				x_1	+	x_2	+	s_1		= 6
2			-	x_1	+	$2x_2$			+	s_2 = 8

The Simplex Algorithm

Canonical Form 0:

Row	Basic Variable							RHS
0	$z=0$	z	$-$	x_1	$-$	$3x_2$		$= 0$
1	$s_1=6$			x_1	$+$	x_2	$+$	$s_1 = 6$
2	$s_2=8$		$-$	x_1	$+$	$2x_2$	$+$	$s_2 = 8$

Is this current basic feasible solution optimal?

To answer this question, we should determine whether there is any way that z can be increased by increasing some nonbasic variable from its current value of zero while holding all other nonbasic variables at their current values of zero.

The Simplex Algorithm

Canonical Form 0:

Row	Basic Variable									RHS
0	$z=0$	z	-	x_1	-	$3x_2$				= 0
1	$s_1=6$			x_1	+	x_2	+	s_1		= 6
2	$s_2=8$		-	x_1	+	$2x_2$			+	s_2 = 8

The Simplex Algorithm

Canonical Form 0:

Row	Basic Variable							RHS
0	$z=0$	z	$-$	x_1	$-$	$3x_2$		$= 0$
1	$s_1=6$			x_1	$+$	x_2	$+$	$s_1 = 6$
2	$s_2=8$		$-$	x_1	$+$	$2x_2$	$+$	$s_2 = 8$

Row 0 : $z - x_1 - 3x_2 = 0$

- if x_1 is increased by 1 unit, z increases by 1 unit
- if x_2 is increased by 1 unit, z increases by 3 units

So we choose x_2 as the **“entering variable”**. If x_2 is to increase from its current value of zero, it has to become a **basic variable**.

The Simplex Algorithm

- For a max. problem, the entering variable has a negative coefficient in row 0. Usually we choose the variable with the most negative coefficient to be the entering variable (ties may be broken in an arbitrary fashion).
- x_2 will become basic. Therefore, one basic variable should become nonbasic. This will be the **“leaving variable”**.

The Simplex Algorithm

- Increasing x_2 may cause a basic variable to become negative. We look at how increasing x_2 (while holding $x_1=0$) changes the values of current set of basic variables:

$$\begin{array}{rcll} \underbrace{x_1}_{=0} + x_2 + s_1 & = & 6 & \\ -\underbrace{x_1}_{=0} + 2x_2 + s_2 & = & 8 & \end{array} \quad \Rightarrow \quad \begin{array}{rcl} x_2 + s_1 & = & 6 \\ 2x_2 + s_2 & = & 8 \end{array}$$

The Simplex Algorithm

$$\text{As } s_1 \geq 0, \quad s_1 = 6 - x_2 \geq 0 \quad \Rightarrow \quad x_2 \leq 6$$

$$\text{As } s_2 \geq 0, \quad s_2 = 8 - 2x_2 \geq 0 \quad \Rightarrow \quad x_2 \leq 4$$

So, x_2 can be at most 4 (otherwise s_2 would become negative!)

The Simplex Algorithm

Observe that for any row in which the entering variable has a positive coefficient, the row's basic variable becomes negative if the entering variable exceeds:

$$\frac{\text{Right Hand Side of row}}{\text{Coefficient of entering variable in row}}$$

The Simplex Algorithm

Ratio Test: When entering a variable into the basis, for every row i in which the entering variable has a positive coefficient, we compute the ratio:

$$\frac{\text{RHS of row } i}{\text{Coefficient of entering variable in row } i},$$

and determine the smallest one.

The smallest ratio is the largest value of the entering variable that will keep all the current basic variables nonnegative.

The Simplex Algorithm

Canonical Form 0:

Row	Basic Variable							RHS
0	$z=0$	z	$-$	x_1	$-$	$3x_2$		$= 0$
1	$s_1=6$			x_1	$+$	x_2	$+$	$s_1 = 6$
2	$s_2=8$		$-$	x_1	$+$	$2x_2$	$+$	$s_2 = 8$

If $x_2=4$, then $s_2=0$ and $s_1=2$.

Therefore s_2 is the **leaving variable** and **becomes nonbasic**.

New basic variables = $\{s_1, x_2\}$; and new nonbasic variables = $\{x_1, s_2\}$

Hence, new $z = x_1 + 3x_2 = 12$.

The Simplex Algorithm

Canonical Form 0:

Row	Basic Variable	entering variable ↓						RHS
0	$z=0$	z	$-$	x_1	$-$	$3x_2$		$= 0$
1	$s_1=6$			x_1	$+$	x_2	$+$	$s_1 = 6$
2	s₂ $=8$		$-$	x_1	$+$	$2x_2$	$+$	$s_2 = 8$

↑
leaving variable

Always make the entering variable a basic variable in a row that wins the ratio test (ties may be broken arbitrarily).

The Simplex Algorithm

Canonical Form 0:

Row	Basic Variable	entering variable ↓							RHS			
0	z	z	-	x_1	-	$3x_2$			=	0		
1	$s_1=6$			x_1	+	x_2	+	s_1	=	6		
2	s₂=8		-	x_1	+	$2x_2$			+	s_2	=	8

↑

leaving variable

Always make the entering variable a basic variable in a row that wins the ratio test (ties may be broken arbitrarily).

The Simplex Algorithm

Canonical Form 0:

Row	Basic Variable	entering variable ↓						RHS		
0	z	z	-	x_1	-	$3x_2$		=	0	
1	s_1			x_1	+	x_2	+	s_1	= 6	
2	$s_2=8$ ↑ leaving variable		-	x_1	+	$2x_2$		+	s_2	= 8

Always make the entering variable a basic variable in a row that wins the ratio test (ties may be broken arbitrarily).

The Simplex Algorithm

Row	Basic Variable									RHS
0	z	z	-	x_1	-	$3x_2$				= 0
1	s_1			x_1	+	x_2	+	s_1		= 6
2			-	x_1	+	$2x_2$			+	s_2 = 8

Always make the entering variable a basic variable in a row that wins the ratio test (ties may be broken arbitrarily).

The Simplex Algorithm

Row	Basic Variable									RHS
0	z	z	-	x_1	-	$3x_2$				= 0
1	s_1			x_1	+	x_2	+	s_1		= 6
2	x_2		-	x_1	+	$2x_2$			+	s_2 = 8

↑
entering variable

Always make the entering variable a basic variable in a row that wins the ratio test (ties may be broken arbitrarily).

The Simplex Algorithm

Row	Basic Variable									RHS
0	z	z	-	x_1	-	$3x_2$				= 0
1	s_1			x_1	+	x_2	+	s_1		= 6
2	x_2		-	x_1	+	$2x_2$			+	s_2 = 8

↑
entering variable

Always make the entering variable a basic variable in a row that wins the ratio test (ties may be broken arbitrarily).

To make x_2 a basic variable in row 2, we use elementary row operations to make x_2 has a coefficient of 1 in row 2 and coefficient of 0 in all other rows.

The Simplex Algorithm

Row	Basic Variable									RHS		
0	z	z	-	x_1	-	$3x_2$				=	0	
1	s_1			x_1	+	x_2	+	s_1		=	6	
2	x_2		-	x_1	+	$2x_2$			+	s_2	=	8

↑

entering variable

Always make the entering variable a basic variable in a row that wins the ratio test (ties may be broken arbitrarily).

To make x_1 a basic variable in row 2, we use elementary row operations to make x_1 has a coefficient of 1 in row 2 and coefficient of 0 in all other rows. This procedure is called **pivoting** on row 2.

The Simplex Algorithm

Row	Basic Variable									RHS
0	z	z	-	x_1	-	$3x_2$				= 0
1	s_1			x_1	+	x_2	+	s_1		= 6
2	x_2		-	x_1	+	$2x_2$			+	s_2 = 8

↑
entering variable

Pivoting: Purpose is to rewrite the original problem in an equivalent form where columns corresponding to basic variables form an identity matrix. This allows us to determine the values of entering and leaving variables in the new solution.

The Simplex Algorithm

Row	Basic Variable									RHS		
0	z	z	-	x_1	-	$3x_2$				=	0	
1	s_1			x_1	+	x_2	+	s_1		=	6	
2	x_2		-	x_1	+	$2x_2$			+	s_2	=	8

We may perform elementary row operations step by step, starting from the pivot row, one row at a time.

The Simplex Algorithm

Row	Basic Variable									RHS
0	z	z	-	x_1	-	$3x_2$				= 0
1	s_1			x_1	+	x_2	+	s_1		= 6
2	x_2		-	x_1	+	$2x_2$			+	s_2 = 8

To make x_2 has a coefficient of 1 in row 2:

- multiply row 2 by 0.5

The Simplex Algorithm

Row	Basic Variable									RHS		
0	z	z	-	x_1	-	$3x_2$				=	0	
1	s_1			x_1	+	x_2	+	s_1		=	6	
2	x_2		-	$0.5 x_1$	+	x_2			+	$0.5 s_2$	=	4

To make x_2 has a coefficient of 1 in row 2:

- multiply row 2 by 0.5

The Simplex Algorithm

Row	Basic Variable									RHS		
0	z	z	-	x_1	-	$3x_2$				=	0	
1	s_1			x_1	+	x_2	+	s_1		=	6	
2	x_2		-	$0.5 x_1$	+	x_2			+	$0.5 s_2$	=	4

To make x_2 has a coefficient of 0 in row 1:

- replace row 1 with row 1 – row 2

The Simplex Algorithm

Row	Basic Variable							RHS
0	z	z	-	x_1	-	$3x_2$		= 0
1	s_1			$1.5 x_1$		+	s_1	- $0.5 s_2$ = 2
2	x_2		-	$0.5 x_1$	+	x_2		+ $0.5 s_2$ = 4

To make x_2 has a coefficient of 0 in row 1:

- replace row 1 with row 1 – row 2

The Simplex Algorithm

Row	Basic Variable							RHS
0	z	z	-	x_1	-	$3x_2$		= 0
1	s_1			$1.5 x_1$		+	s_1	- $0.5 s_2$ = 2
2	x_2		-	$0.5 x_1$	+	x_2		+ $0.5 s_2$ = 4

To make x_2 has a coefficient of 0 in row 0:

- replace row 0 with row 0 + 3 (row 2)

The Simplex Algorithm

Row	Basic Variable							RHS
0	z	z	-	2.5 x_1		+	1.5 s_2	= 12
1	s_1			1.5 x_1	+	s_1	- 0.5 s_2	= 2
2	x_2		-	0.5 x_1	+	x_2	+ 0.5 s_2	= 4

To make x_2 has a coefficient of 0 in row 0:

- replace row 0 with row 0 + 3 (row 2)

The Simplex Algorithm

Canonical Form 1:

Row	Basic Variable	RHS									
0	z	z	-	2.5 x_1			+	1.5 s_2	=	12	
1	s_1			1.5 x_1		+	s_1	-	0.5 s_2	=	2
2	x_2		-	0.5 x_1	+	x_2		+	0.5 s_2	=	4

Is this bfs optimal?

No, increasing the nonbasic variable x_1 will increase z !

So x_1 *is the entering variable*. Also note that x_1 is the variable with the most negative coefficient in row 0.

The Simplex Algorithm

We perform the ration test to find the leaving variable:

Since $s_2 = 0$, the system is:

$$1.5 x_1 + s_1 = 2, \text{ and } s_1 \geq 0 \Rightarrow s_1 = 2 - 1.5 x_1 \geq 0$$

$$\Rightarrow x_1 \leq 4/3$$

$$-0.5 x_1 + x_2 = 4, \text{ and } x_2 \geq 0 \Rightarrow x_2 = 4 + 0.5 x_1 \geq 0$$

$$\Rightarrow x_1 \geq -8$$

So, $x_1 = 4/3$ and s_1 becomes the leaving variable

The Simplex Algorithm

The new bfs is:

$$\begin{aligned}x_1 &= 4/3, \\x_2 &= 14/3, \\s_1 &= 0, \\s_2 &= 0, \\z &= 46/3.\end{aligned}$$

Now, keep the above results in mind and let us have a look at the pivot of the simplex algorithm:

The Simplex Algorithm

Canonical Form 1:

Row	Basic Variable	entering variable ↓						RHS
0	z	z	-	2.5 x ₁			+ 1.5 s ₂	= 12
1	s ₁			1.5 x ₁	+	s ₁	- 0.5 s ₂	= 2
2	x ₂		-	0.5 x ₁	+	x ₂	+ 0.5 s ₂	= 4

leaving variable

The Simplex Algorithm

Row	Basic Variable	entering variable ↓						RHS
0	z	z	-	2.5 x ₁		+	1.5 s ₂	= 12
1				1.5 x ₁	+	s ₁	- 0.5 s ₂	= 2
2	x ₂		-	0.5 x ₁	+	x ₂	+ 0.5 s ₂	= 4

↑
leaving variable

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable							RHS
0	z	z	-	2.5 x_1		+	1.5 s_2	= 12
1	x_1			1.5 x_1	+	s_1	- 0.5 s_2	= 2
2	x_2		-	0.5 x_1	+	x_2	+ 0.5 s_2	= 4

To make x_1 a basic variable in row 1, we use elementary row operations to make x_1 has a coefficient of 1 in row 1 and coefficient of 0 in all other rows.

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable							RHS
0	z	z	-	2.5 x_1		+	1.5 s_2	= 12
1	x_1			1.5 x_1	+	s_1	- 0.5 s_2	= 2
2	x_2		-	0.5 x_1	+	x_2	+ 0.5 s_2	= 4

To make x_1 has a coefficient of 1 in row 1:

- multiply row 1 by 2/3

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable							RHS
0	z	z	-	2.5 x_1		+	1.5 s_2	= 12
1	x_1			x_1	+	$\frac{2}{3} s_1$	- $\frac{1}{3} s_2$	= $\frac{4}{3}$
2	x_2		-	0.5 x_1	+	x_2	+	0.5 s_2 = 4

To make x_1 has a coefficient of 1 in row 1:

- multiply row 1 by $\frac{2}{3}$

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable							RHS
0	z	z	-	2.5 x ₁		+	1.5 s ₂	= 12
1	x ₁			x ₁	+	2/3 s ₁	- 1/3 s ₂	= 4/3
2	x ₂		-	0.5 x ₁	+	x ₂	+	0.5 s ₂ = 4

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable							RHS
0	z	z	-	2.5 x_1		+	1.5 s_2	= 12
1	x_1			x_1	+	$\frac{2}{3} s_1$	- $\frac{1}{3} s_2$	= $\frac{4}{3}$
2	x_2		-	0.5 x_1	+	x_2	+	0.5 s_2 = 4

To make x_1 has a coefficient of 0 in row 2:

- multiply row 1 by 0.5 and add to the row 2

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable						RHS
0	z	z	-	2.5 x_1		+ 1.5 s_2	= 12
1	x_1			x_1	+ $\frac{2}{3} s_1$	- $\frac{1}{3} s_2$	= $\frac{4}{3}$
2	x_2			x_2	+ $\frac{1}{3} s_1$	+ $\frac{1}{3} s_2$	= $\frac{14}{3}$

To make x_1 has a coefficient of 0 in row 2:

- multiply row 1 by 0.5 and add to the row 2

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable							RHS
0	z	z	-	2.5 x_1		+	1.5 s_2	= 12
1	x_1			x_1	+	$\frac{2}{3} s_1$	- $\frac{1}{3} s_2$	= $\frac{4}{3}$
2	x_2			x_2	+	$\frac{1}{3} s_1$	+ $\frac{1}{3} s_2$	= $\frac{14}{3}$

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable						RHS
0	z	z	-	2.5 x_1		+ 1.5 s_2	= 12
1	x_1			x_1	+ $\frac{2}{3} s_1$	- $\frac{1}{3} s_2$	= $\frac{4}{3}$
2	x_2			x_2	+ $\frac{1}{3} s_1$	+ $\frac{1}{3} s_2$	= $\frac{14}{3}$

To make x_1 has a coefficient of 0 in row 0:

- multiply row 1 by 2.5 and add to the row 0

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable						RHS
0	z	z		+ 5/3 s ₁	+ 2/3 s ₂	=	46/3
1	x ₁		x ₁	+ 2/3 s ₁	- 1/3 s ₂	=	4/3
2	x ₂			x ₂	+ 1/3 s ₁	+ 1/3 s ₂	= 14/3

To make x₁ has a coefficient of 0 in row 0:

- multiply row 1 by 2.5 and add to the row 0

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable						RHS
0	z	z		+ 5/3 s ₁	+ 2/3 s ₂	=	46/3
1	x ₁		x ₁	+ 2/3 s ₁	- 1/3 s ₂	=	4/3
2	x ₂			x ₂	+ 1/3 s ₁	+ 1/3 s ₂	= 14/3

This result is the same as we had calculated before!

The Simplex Algorithm

Previously calculated new bfs was:

$$x_1 = 4/3,$$

$$x_2 = 14/3,$$

$$s_1 = 0,$$

$$s_2 = 0,$$

$$z = 46/3.$$

The Simplex Algorithm

Canonical Form 2:

Row	Basic Variable						RHS
0	z	z		+ 5/3 s ₁	+ 2/3 s ₂	=	46/3
1	x ₁		x ₁	+ 2/3 s ₁	- 1/3 s ₂	=	4/3
2	x ₂			x ₂	+ 1/3 s ₁	+ 1/3 s ₂	= 14/3

Is this bfs optimal?

YES! Because increasing nonbasic variables s_1 and s_2 will decrease z (Also note that there is no variable in row 0 with a negative coefficient!)

Representing Simplex Tableaus

- Instead of writing each variable in every constraint, we can use a shorthand display called a ***simplex tableau***.

Representing Simplex Tableaus

Canonical Form 0:

Row	Basic Variable									RHS		
0	z=0	z	-	x ₁	-	3x ₂				=	0	
1	s ₁ =6			x ₁	+	x ₂	+	s ₁		=	6	
2	s ₂ =8		-	x ₁	+	2x ₂			+	s ₂	=	8

For example, canonical form 0 could be written as

Row	Basic Variable	z	x_1	x_2	s_1	s_2	RHS
0	$z=0$	1	-1	-3	0	0	0
1	$s_1=6$	0	1	1	1	0	6
2	$s_2=8$	0	-1	2	0	1	8

Representing Simplex Tableaus

- With this format, it is very easy to spot the basic variables. We just look for columns with entry of 1 and all entries equal to 0.

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z=0	1	-1	-3	0	0	0
$s_1=6$	0	1	1	1	0	6
$s_2=8$	0	-1	2	0	1	8

Note that z will always be a basic variable. Therefore, we won't be mentioning it unless it is necessary. In addition, since row 0 corresponds to the objective function, it is indicated separately in the simplex tableau.

Also note that, since we are in canonical form, the basic variable of a row will be equal to the RHS of that row.

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z=0	1	-1	-3	0	0	0
$s_1=6$	0	1	1	1	0	6
$s_2=8$	0	-1	2	0	1	8

Now let us summarize what we have done so far!

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
	0	1	1	1	0	6
	0	-1	2	0	1	8

For a max problem:

We start with an initial bfs in the canonical form above (if there are m slack variables, we use them as basic variables)

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
s_1	0	1	1	1	0	6
s_2	0	-1	2	0	1	8

For a max problem:

We start with an initial bfs in the canonical form above (if there are m slack variables, we use them as basic variables)

Representing Simplex Tableaus

Simplex Tableau-0: entering variable



Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
s_1	0	1	1	1	0	6
s_2	0	-1	2	0	1	8

Entering Variable: Choose a variable with the most negative coefficient in row 0.

Representing Simplex Tableaus

Simplex Tableau-0: entering variable



Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
s_1	0	1	1	1	0	6
s_2	0	-1	2	0	1	8

Representing Simplex Tableaus

Simplex Tableau-0: entering variable



Basic Variable	z	x ₁	x ₂	s ₁	s ₂	RHS
z	1	-1	-3	0	0	0
s ₁	0	1	1	1	0	6
s ₂	0	-1	2	0	1	8

Leaving Variable: compute ratios:

$$\frac{\text{RHS of row } i}{\text{Coefficient of entering variable in row } i'}$$

The smallest positive ratio wins and the basic variable of the winning row leaves the basis.

Representing Simplex Tableaus

Simplex Tableau-0:

entering variable

Basic Variable	z	x_1	x_2	s_1	s_2	RHS	Ratio Test
z	1	-1	-3	0	0	0	
s_1	0	1	1	1	0	6	6
s_2	0	-1	2	0	1	8	4

leaving variable

pivot term

Leaving Variable: compute ratios:

$$\frac{\text{RHS of row } i}{\text{Coefficient of entering variable in row } i'}$$

The smallest positive ratio wins and the basic variable of the winning row leaves the basis.

Representing Simplex Tableaus

Simplex Tableau-0:

entering variable

Basic Variable	z	x_1	x_2	s_1	s_2	RHS	Ratio Test
z	1	-1	-3	0	0	0	
s_1	0	1	1	1	0	6	6
s_2	0	-1	2	0	1	8	4

leaving variable

pivot term

Leaving Variable: compute ratios:


$$\frac{\text{RHS of row } i}{\text{Coefficient of entering variable in row } i'}$$

The smallest positive ratio wins and the basic variable of the winning row leaves the basis.

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS	Ratio Test
z	1	-1	-3	0	0	0	
s_1	0	1	1	1	0	6	6
x_2	0	-1	2	0	1	8	4



Leaving Variable: compute ratios:


$$\frac{\text{RHS of row } i}{\text{Coefficient of entering variable in row } i'}$$

The smallest positive ratio wins and the basic variable of the winning row leaves the basis.

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
s_1	0	1	1	1	0	6
x_2	0	-1	2	0	1	8



Leaving Variable: compute ratios:


$$\frac{\text{RHS of row } i}{\text{Coefficient of entering variable in row } i'}$$

The smallest positive ratio wins and the basic variable of the winning row leaves the basis.

Representing Simplex Tableaus

Simplex Tableau-0:


Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
s_1	0	1	1	1	0	6
x_2	0	-1	2	0	1	8



Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
s_1	0	1	1	1	0	6
x_2	0	-1	2	0	1	8




Pivot: Transform the tableau so that the new basic variable (entering variable) has 1 in the row of the leaving variable (pivot row) and 0 in other rows.

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
s_1	0	1	1	1	0	6
x_2						




Pivot: Transform the tableau so that the new basic variable (entering variable) has 1 in the row of the leaving variable (pivot row) and 0 in other rows.

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
s_1	0	1	1	1	0	6
x_2	0	-1/2	1	0	1/2	4




Pivot: Transform the tableau so that the new basic variable (entering variable) has 1 in the row of the leaving variable (pivot row) and 0 in other rows.

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
s_1	0	-1/2	1	0	1/2	4
x_2						




Pivot: Transform the tableau so that the new basic variable (entering variable) has 1 in the row of the leaving variable (pivot row) and 0 in other rows.

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	-1	-3	0	0	0
s_1	0	$3/2$	0	1	$-1/2$	2
x_2	0	$-1/2$	1	0	$1/2$	4




Pivot: Transform the tableau so that the new basic variable (entering variable) has 1 in the row of the leaving variable (pivot row) and 0 in other rows.

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z						
s_1	0	$3/2$	0	1	$-1/2$	2
x_2	0	$-1/2$	1	0	$1/2$	4




Pivot: Transform the tableau so that the new basic variable (entering variable) has 1 in the row of the leaving variable (pivot row) and 0 in other rows.

Representing Simplex Tableaus

Simplex Tableau-0:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	$-5/2$	0	0	$3/2$	12
s_1	0	$3/2$	0	1	$-1/2$	2
x_2	0	$-1/2$	1	0	$1/2$	4



Pivot: Transform the tableau so that the new basic variable (entering variable) has 1 in the row of the leaving variable (pivot row) and 0 in other rows.

Representing Simplex Tableaus

Simplex Tableau-1:

Basic Variable	z	enter ↓ x_1	x_2	s_1	s_2	RHS
z	1	-5/2	0	0	3/2	12
s_1	0	3/2	0	1	-1/2	2
x_2	0	-1/2	1	0	1/2	4

Representing Simplex Tableaus

Simplex Tableau-1:

Basic Variable	z	enter				RHS	Ratio Test
		x_1	x_2	s_1	s_2		
z	1	-5/2	0	0	3/2	12	
← s_1	0	3/2	0	1	-1/2	2	4/3
x_2	0	-1/2	1	0	1/2	4	no ratio

Representing Simplex Tableaus

Simplex Tableau-1:

Basic Variable	z	enter				RHS	Ratio Test
		x_1	x_2	s_1	s_2		
z	1	-5/2	0	0	3/2	12	
leave ←	0	3/2	0	1	-1/2	2	4/3
x_2	0	-1/2	1	0	1/2	4	no ratio

Representing Simplex Tableaus

Simplex Tableau-1:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS	Ratio Test
z	1	$-5/2$	0	0	$3/2$	12	
x_1	0	$3/2$	0	1	$-1/2$	2	4/3
x_2	0	$-1/2$	1	0	$1/2$	4	no ratio

Representing Simplex Tableaus

Simplex Tableau-1:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	$-5/2$	0	0	$3/2$	12
x_1	0	$3/2$	0	1	$-1/2$	2
x_2	0	$-1/2$	1	0	$1/2$	4

Representing Simplex Tableaus

Simplex Tableau-2:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	0	0	$5/3$	$2/3$	$46/3$
x_1	0	1	0	$2/3$	$-1/3$	$4/3$
x_2	0	0	1	$1/3$	$1/3$	$14/3$

Representing Simplex Tableaus

Simplex Tableau-2:

Basic Variable	z	x_1	x_2	s_1	s_2	RHS
z	1	0	0	$5/3$	$2/3$	$46/3$
x_1	0	1	0	$2/3$	$-1/3$	$4/3$
x_2	0	0	1	$1/3$	$1/3$	$14/3$

Can we iterate more?

No, because all row 0 coefficients are nonnegative. We stop here. The current bfs is optimal!

Summary of the Simplex Algorithm

(For a max problem)

- 1) Convert the LP into the standard form and then obtain the canonical form
- 2) Find an initial bfs (if there are m slack variables, use them as basic variables).

If all nonbasic variables have nonnegative coefficients in row 0, then the current LP is optimal.

If there are any variables with a negative coefficient, then we should decide the entering variable.

Entering variable: choose a variable with the most negative coefficient in row 0 to enter the basis.

Summary of the Simplex Algorithm

- 3) For any row in which the entering variable has a *positive coefficient*, compute the ratios:

$$\frac{\text{RHS of row } i}{\text{Coefficient of the entering variable in row } i}$$

The smallest ratio wins (ties may be broken arbitrarily) and the basic variable of the winning row leaves the basis.

- 4) **Pivot:** Transform the tableau so that the new basic variable (entering variable) has coefficient of 1 in the row of the leaving variable (pivot row) and coefficient of 0 in all other rows. In the end, we get a tableau with a new canonical form.

Summary of the Simplex Algorithm

After the pivoting, note that:

$$\text{New pivot row} = \frac{1}{\left(\begin{array}{c} \text{coefficient of the entering} \\ \text{variable in the pivot row} \end{array} \right)} \cdot (\text{old pivot row})$$

$$\text{new row } i = \text{old row } i - \left(\begin{array}{c} \text{coefficient of the entering} \\ \text{variable in the pivot row} \end{array} \right) \cdot (\text{new pivot row})$$

Summary of the Simplex Algorithm

5) **Repeat steps 1,2,3, and 4** until all row 0 coefficients becomes nonnegative. If each nonbasic variable has a nonnegative coefficient in a canonical form's row 0 (remember that basic variables have coefficient of 0 in row 0 of a canonical form), then the canonical form is optimal. ***We stop here and the current bfs is optimal!***

